arXiv:0705.4127v1 [math.AG] 29 May 2007

THE AUTOMORPHISM GROUP OF TORIC DELIGNE-MUMFORD STACKS

YUNFENG JIANG

Abstract. We prove that the automorphism group of a toric Deligne-Mumford stack is isomorphic to the 2-group associated to the stacky fan.

1. Introduction

Let Σ be a simplicial fan with n rays in a lattice N of dimension d. The simplicial toric variety $X(\Sigma)$ can be represented as a geometric quotient Z/G, where $Z := \mathbb{C}^n \setminus V(J_\Sigma)$, $V(J_\Sigma)$ is the closed subvariety defined by the irrelevant ideal J_Σ of the fan Σ, see [Cox]. The group $G = \mathrm{Hom}(A_{d-1}(X(\Sigma)), \mathbb{C}^*)$ acts on Z through the map $\alpha : G \to (\mathbb{C}^*)^n$ obtained by taking $\mathrm{Hom}(-, \mathbb{C}^*)$-functor on the map $\beta^\vee$ in the following exact sequence

$$0 \longrightarrow N^\star \longrightarrow \mathbb{Z}^n \xrightarrow{\beta^\vee} A_{d-1}(X(\Sigma)) \longrightarrow 0, \tag{1.1}$$

in [F], where $A_{d-1}(X(\Sigma))$ is the Weil divisor class group of the toric variety $X(\Sigma)$. Let $\mathrm{Aut}(X(\Sigma))$ be the automorphism group of $X(\Sigma)$ and $\mathrm{Aut}(Z)$ the automorphism group of Z. Since the group G acts on Z, it is naturally a subgroup of the automorphism group $\mathrm{Aut}(Z)$. Let $\widetilde{\mathrm{Aut}(Z)}$ be the normalizer of G in $\mathrm{Aut}(Z)$. Then from [Cox] we have the following exact sequence

$$1 \longrightarrow G \longrightarrow \widetilde{\mathrm{Aut}(Z)} \longrightarrow \mathrm{Aut}(X(\Sigma)) \longrightarrow 1. \tag{1.2}$$

If $\widetilde{\mathrm{Aut}^0(Z)}$ and $\mathrm{Aut}^0(X(\Sigma))$ are the components of the identity elements in $\widetilde{\mathrm{Aut}(Z)}$ and $\mathrm{Aut}(X(\Sigma))$, then we have the exact sequence

$$1 \longrightarrow G \longrightarrow \widetilde{\mathrm{Aut}^0(Z)} \longrightarrow \mathrm{Aut}^0(X(\Sigma)) \longrightarrow 1. \tag{1.3}$$

The main goal of this note is to generalize the above results to toric Deligne-Mumfords stacks. Generalizing the idea of Cox on simplicial toric varieties, Borisov, Chen and Smith [BCS] defined the notion of toric Deligne-Mumford stacks which are encoded in terms of stacky fans. A stacky fan $\mathbf{\Sigma} = (N, \Sigma, \beta)$ is a triple, where N is a finitely generated abelian group, $\Sigma \subset N \otimes_{\mathbb{Z}} \mathbb{Q}$ is a simplicial fan and $\beta : \mathbb{Z}^n \longrightarrow N$ is a map determined by

Date: February 1, 2008.

the elements $\{b_1, \cdots, b_n\}$ in N. We require that β has finite cokernel and $\{\overline{b}_1, \cdots, \overline{b}_n\}$ generate the simplicial fan Σ, where $\overline{b}_i$ is the image of b_i under the natural map $N \longrightarrow \overline{N} = N/N_{tor}$. Then we have the exact sequences:

$$0 \longrightarrow DG(\beta)^\star \longrightarrow \mathbb{Z}^n \xrightarrow{\beta} N \longrightarrow Coker(\beta) \longrightarrow 0, \tag{1.4}$$

and

$$0 \longrightarrow N^\star \longrightarrow \mathbb{Z}^n \xrightarrow{\beta^\vee} DG(\beta) \longrightarrow Coker(\beta^\vee) \longrightarrow 0, \tag{1.5}$$

where $\beta^\vee$ is the Gale dual of β (see [BCS]). The toric Deligne-Mumford stack $\mathcal{X}(\mathbf{\Sigma})$ associated to $\mathbf{\Sigma}$ is defined to be the quotient stack $[Z/G]$, where Z is the same as in the quotient construction of toric varieties, $G = \mathrm{Hom}(DG(\beta), \mathbb{C}^*)$ and the action is through a group homomorphism $\alpha : G \longrightarrow (\mathbb{C}^*)^n$ in the exact sequence

$$1 \longrightarrow \mu \longrightarrow G \xrightarrow{\alpha} (\mathbb{C}^*)^n \longrightarrow (\mathbb{C}^*)^d \longrightarrow 1, \tag{1.6}$$

which is obtained by taking $\mathrm{Hom}(-, \mathbb{C}^*)$-functor on the exact sequence (1.5).

Let $\mathcal{X}$ be an algebraic stack. The automorphism group of $\mathcal{X}$ is naturally a 2-group because an isomorphism of the stack is a 2-isomorphism. A 2-group is a group object in the category of groupoids. The notion of 2-groups, also called crossed-modules, appeared first in algebraic topology. The fundamental concepts of 2-groups and crossed-models can be found in [Noohi]. A crossed-module $\mathfrak{G} = [G_2 \to G_1]$ consists of a pair of groups G_1 and G_2, a group homomorphism $\varphi : G_2 \to G_1$, and a (right) action of G_1 on G_2, denoted by ${}^{-}a$, which lifts the conjugation action of G_1 on the image of φ and descends the conjugation action of G_2 on itself.

Let $\mathcal{X}(\mathbf{\Sigma})$ be a toric Deligne-Mumford stack associated to the stacky fan $\mathbf{\Sigma}$. Since $(\mathbb{C}^*)^n$ is the maximal torus of the automorphism group $\mathrm{Aut}(Z)$, the map α in the exact sequence (1.6) naturally induces a map $\varphi : G \to \widetilde{\mathrm{Aut}(Z)}$ whose image lies in the centralizer $\widetilde{\mathrm{Aut}^0(Z)}$ of G in $\widetilde{\mathrm{Aut}(Z)}$. We define the action of $\widetilde{\mathrm{Aut}(Z)}$ on G by:

$$\begin{cases} h \cdot g = g & \text{if } \ h \in \widetilde{\mathrm{Aut}^0(Z)}; \\ \\ h \cdot g = hgh^{-1} & \text{if } h \in \widetilde{\mathrm{Aut}(Z)} \setminus \widetilde{\mathrm{Aut}^0(Z)}\,. \end{cases} \tag{1.7}$$

Then

$$\mathfrak{G} = [G \xrightarrow{\varphi} \widetilde{\mathrm{Aut}(Z)}]$$

is a crossed module and it defines a (weak) 2-group. We call $\mathfrak{G}$ the 2-group associated to the stacky fan $\mathbf{\Sigma}$. Let $\mathrm{Aut}(\mathcal{X}(\mathbf{\Sigma}))$ be the (weak) 2-group of the automorphism group of the toric Deligne-Mumford stack $\mathcal{X}(\mathbf{\Sigma})$. Then there is a natural homomorphism from $\mathfrak{G}$ to $\mathrm{Aut}(\mathcal{X}(\mathbf{\Sigma}))$ which will be discussed in Section 2.

Theorem 1.1. *Let $\mathcal{X}(\mathbf{\Sigma})$ be a toric Deligne-Mumford stack associated to the stacky fan $\mathbf{\Sigma}$. Then the natural map*

$$f : \mathfrak{G} \to Aut(\mathcal{X}(\mathbf{\Sigma}))$$

is an isomorphism.

Actually we prove a result for general quotient stack $[X/G]$ in which the group G is abelian such that the theorem is a corollary of this result. Our result generalizes Lemma 8.2 of Behrend and Noohi [BN],where they only consider the case of the centralizer of the group G in the automorphism group of X.

The paper is outlined as follows. In Section 2 we quickly review the notion of 2-groups. The main Theorem is proved in Section 3. In Section 4 we prove that the weighted projective stacks are toric Deligne-Mumford stacks and discuss the weighted projective linear 2-groups which are the automorphism group of weighted projective stacks.

Conventions. In this paper we work entirely algebraically over the field of complex numbers. By an orbifold we mean a smooth Deligne-Mumford stack with trivial generic stabilizer.

We write $N^\star$ for $Hom_{\mathbb{Z}}(N, \mathbb{Z})$ and $N \to \overline{N}$ the natural map of modulo torsion. We refer the reader to [BCS] for the construction of the Gale dual $\beta^\vee : \mathbb{Z}^n \to DG(\beta)$ of $\beta : \mathbb{Z}^n \to N$.

We use the notation $\mathfrak{G}$ to represent 2-groups. For more details about 2-groups and crossed modules, see [Noohi].

Acknowledgments. I would like to thank my advisor Kai Behrend for encouragements and valuable discussions.

2. 2-groups and automorphism group of stacks

In this section we review the notions of 2-groups and crossed modules. The automorphism group of an algebraic stack is naturally a 2-group.

2.1. **2-groups.** A 2-group $\mathfrak{G}$ is a group object in the category of groupoids. Alternatively, we can define a 2-group to be a groupoid object in the category of groups, or also, as a (strict) 2-category with one object in which all 1-morphisms and 2-morphisms are invertible (in the strict sense). If we require that the 1-morphisms are only equivalences and not necessarily strictly invertible, what we obtain is called a weak 2-group.

A morphism $f : \mathfrak{G} \to \mathfrak{H}$ of 2-groups is a map of groupoids that respects the group operation on the nose. If we view $\mathfrak{G}$ and $\mathfrak{H}$ as 2-categories with one object, such f is nothing but a strict 2-functor. Same definition applies to

weak 2-groups (because we have not weakened associativity). To a 2-group $\mathfrak{G}$ we associate the groups $\pi_1(\mathfrak{G})$ and $\pi_2(\mathfrak{G})$ as follows. The group $\pi_1(\mathfrak{G})$ is the set of isomorphism classes of object of the groupoid $\mathfrak{G}$. The group structure on $\pi_1(\mathfrak{G})$ is induced from the group operation of $\mathfrak{G}$. The group $\pi_2(\mathfrak{G})$ is the group of automorphisms of the identity object e in $\mathfrak{G}$. This is an abelian group. Any morphism $f : \mathfrak{G} \to \mathfrak{H}$ of (weak) 2-groups induces homomorphisms on $\pi_1(\mathfrak{G})$ and $\pi_2(\mathfrak{G})$. We say such f is an *equivalence* if both these maps are isomorphisms. (Warning: an equivalence need not have an inverse. Also, two equivalent 2-groups may not be related by an equivalence, but just a zig-zag of equivalences.) We are usually interested in 2-groups up to equivalence, so we will think of a 2-group as an object in the homotopy category of 2-groups. This category is defined by taking the category of 2-groups and inverting the equivalences. For the readers information, we point out that, there is a model structure on the category of 2-groupoids; 2-groups are the pointed connected objects in this category.

2.2. **Crossed moduels.** A crossed-module $\mathfrak{G} = [\varphi : G_2 \to G_1]$ consists of a pair of groups G_1 and G_2, a group homomorphism $\varphi : G_2 \to G_1$, and a (right) action of G_1 on G_2, denoted by ${}^{-}g$, that lifts the conjugation action of G_1 on the image of φ and descends the conjugation action of G_2 on itself. The kernel of φ is a central (in particular abelian) subgroup of G_2 and is denoted by $\pi_2(\mathfrak{G})$. The image of φ is a normal subgroup of G_1 whose cokernel is denoted by $\pi_1(\mathfrak{G})$. A (strict) morphism of crossed-modules is a pair of group homomorphisms which commute with the φ maps and respect the action of G_1 on G_2. A morphism is called an equivalence if it induces an isomorphism on $\pi_1(\mathfrak{G})$ and $\pi_2(\mathfrak{G})$.

Equivalence of 2-groups and crossed-modules. There is a natural pair of inverse equivalences of categories between the category **2Gp** of 2-groups and the category **CrossedMod** of crossed-modules. Furthermore, these functors preserve $\pi_1(\mathfrak{G})$ and $\pi_2(\mathfrak{G})$. Here is how these functors are defined.

Functor from 2-groups to crossed-modules. Let $\mathfrak{G}$ be a 2-group. Let G_1 be the group of objects of $\mathfrak{G}$, and let G_2 be the set of all arrows emanating from the identity object e; G_2 is also a group (namely, it is a subgroup of the group of arrows of $\mathfrak{G}$). Define

$$\varphi : G_2 \to G_1$$

by $\varphi(\alpha) := t(\alpha)$, for $\alpha \in G_2$. The action of G_1 on G_2 is given by conjugation. That is, given $\alpha \in G_2$ and $g \in G_1$, the action is given by $g^{-1}\alpha g$.

Functor from crossed-modules to 2-groups. Let $[\varphi : G_2 \to G_1]$ be a crossed-module. Consider the groupoid $\mathfrak{G}$ whose underlying set of objects is G_1 and whose set of arrows is $G_1 \times G_2$. The source and target maps are given by $s(g, \alpha) = g$, $t(g, \alpha) = g\varphi(\alpha)$. Two arrows (g, α) and (h, β) such

that $g\varphi(\alpha) = h$ are composed to $(g, \alpha\beta)$. The group operation on the set of objects $\mathbf{ObG} = G_1$ is naturally extended to a group operation on $\mathfrak{G}$ by setting $(g, \alpha)(h, \beta) = (gh, (\alpha^h)\beta)$, where $-^h$ stands for the action of G_1 on G_2.

2.3. **Automorphism group of stacks.** Let $\mathcal{X}$ be a smooth Deligne-Mumford stack. From [L-MB], let $X \longrightarrow \mathcal{X}$ be an etale presentation of the stack $\mathcal{X}$, then $\mathcal{X}$ can be represented by a etale groupoid $\mathcal{X} = [X \times_{\mathcal{X}} X \rightrightarrows X]$ with source and target maps s, t. For the groupoid $\mathcal{X}$, the automorphism group $\mathfrak{G} = \mathrm{Aut}(\mathcal{X})$ is a (weak) 2-group. We have the associated groups $\pi_1(\mathfrak{G})$ and $\pi_2(\mathfrak{G})$ which are defined in Section 2.1.

We are mainly interested in the quotient stack $\mathcal{X} = [X/G]$ in which the group G is an abelian group. So the associated groupoid is

$$X \times G \rightrightarrows X.$$

Let $\mathrm{Aut}(X)$ be the automorphism group of X, then we have an embedding $G \longrightarrow \mathrm{Aut}(X)$. Let $\widetilde{\mathrm{Aut}(X)}$ be the normalizer of G in $\mathrm{Aut}(X)$ and $\widetilde{\mathrm{Aut}^0(X)}$ the centralizer of G in $\mathrm{Aut}(X)$, i.e. the component of the identity element in $\widetilde{\mathrm{Aut}(X)}$. We define the action of $\widetilde{\mathrm{Aut}(X)}$ on G as follows: for $g \in G$,

$$\begin{cases} h \cdot g = g & \text{if } \ h \in \widetilde{\mathrm{Aut}^0(X)}; \\ \\ h \cdot g = hgh^{-1} & \text{if } h \in \widetilde{\mathrm{Aut}(X)} \setminus \widetilde{\mathrm{Aut}^0(X)}\,; \end{cases} \tag{2.1}$$

Then we have a 2-group

$$\mathfrak{G} := [G \to \widetilde{\mathrm{Aut}(X)}].$$

From [BN], there is a natural morphism

$$f : \mathfrak{G} \longrightarrow \mathrm{Aut}(\mathcal{X}).$$

In Section 3, we prove that this morphism φ is an isomorphism.

3. The proof of the Theorem

In this section we prove the main theorem of this note. Our proof bases on a generalization of Lemma 8.2 of Behrend and Noohi [BN] such that our main result is a corollary of the generalization. We first recall the key lemma of Behrend and Noohi for the automorphism group of abelian quotient stacks.

Lemma 3.1. *(*[BN]*) Let G be an abelian group scheme acting on a connected scheme X over $\mathbb{C}$, and let $\mathcal{X} = [X/G]$ be the quotient stack. Let $\widetilde{Aut^0(X)}$ be the centralizer of G in the automorphism group $Aut(X)$ and $Aut(\mathcal{X})$ the (weak) 2-group of automorphisms of $\mathcal{X}$. Then*

(1) *The natural homomorphism* $\varphi : G \to \widetilde{Aut^0(X)}$ *can be turned into a crossed moduel by taking the trivial action of* $\widetilde{Aut^0(X)}$ *on* G*;*
(2) *If* $\mathfrak{G} = [G \xrightarrow{\varphi} \widetilde{Aut^0(X)}]$ *is the* 2*-group associated to the crossed moduel. Then there is a natural map of* 2*-groups* $\mathfrak{G} \to Aut(\mathcal{X})$*. Furthermore, this morphism induces an isomorphism on* π_2*;*
(3) *Assume that* $\mathcal{X}$ *is a proper Deligne-Mumford stack and* G *is affine. Then the induced map on* π_1 *is injective.* □

We construct a new crossed moduel from the quotient stack so that the associated 2-group is equivalent to the (weak) automorphism group of the quotient stack.

Lemma 3.2. *Let* G *be an abelian group scheme acting on a connected scheme* X *over* $\mathbb{C}$*, and let* $\mathcal{X} = [X/G]$ *be the quotient stack. Let* $\widetilde{Aut(X)}$ *be the normalizer of* G *in the automorphism group* $Aut(X)$ *and* $Aut(\mathcal{X})$ *the (weak)* 2*-group of automorphisms of* $\mathcal{X}$*. Then*

(1) *The natural homomorphism* $\varphi : G \to \widetilde{Aut(X)}$ *can be turned into a crossed moduel by taking the action (2.1) of* $\widetilde{Aut(X)}$ *on* G*.*
(2) *If* $\mathfrak{G} = [G \xrightarrow{\varphi} \widetilde{Aut(X)}]$ *is the* 2*-group associated to the crossed moduel. Then the natural map of* 2*-groups* $f : \mathfrak{G} \to Aut(\mathcal{X})$ *is an equivalence as* 2*-groups.*

Proof.

(1) Since φ maps G to the center and the given action of $\widetilde{\mathrm{Aut}(X)}$ on G extends the action of G on itself by conjugation to $\widetilde{\mathrm{Aut}(X)}$, from the definition of crossed moduel, $\mathfrak{G}$ is a crossed moduel which corresponds to a 2-group;
(2) To prove that the natural map $f : \mathfrak{G} \to \mathrm{Aut}(\mathcal{X})$ is an isomorphism, it suffices to prove that the map induces isomorphisms on π_2 and π_1. We first consider the π_2 case. We prove that any element $g \in \mathrm{Aut}(\mathcal{X})$ induces an isomorphism of $\mathcal{X}$. Recall that $\mathcal{X} = [X/G]$ is a quotient stack,

$$ob[X/G](S) = \{(T, \alpha) | T \text{ a } G \text{ torsor over } S, \alpha : T \to X \text{ an } G \text{ map}\},$$

and

$$Mor[X/G](S)((T, \alpha), (T', \alpha')) = \{f : T \to T' \text{ an } G \text{ torsor map such that } \alpha' \circ f = \alpha\}.$$

If $g \in \mathrm{Aut}^0(\mathcal{X})$, then the induced automorphism of $\mathcal{X}$ is given by keeping the same torsor T and compose α with the action of g on X which is the same as in [BN]. If $g \in \mathrm{Aut}(\mathcal{X}) \setminus \mathrm{Aut}^0(\mathcal{X})$, the we have

the diagram:

$$\begin{array}{ccc} T & \xrightarrow{f} & T' \\ {\scriptstyle\alpha}\downarrow & \swarrow_{g\circ\alpha} & \\ X, & & \end{array} \tag{3.1}$$

then $g\circ\alpha$ is a new G-torsor $T' \xrightarrow{\alpha'} X$, which we have an isomorphism between T to T'. Hence we can define a morphism of 2-groups

$$f : \mathfrak{G} \to \mathrm{Aut}(\mathcal{X}).$$

The proof of isomorphism on π_2 is the same as in [BN], we omit the details.

We now prove the isomorphism on π_1. The injectivity on π_1 is the same as in [BN]. We prove the surjectivity. Let g be an automorphism of the isomorphism classes of the objects on $\mathcal{X}$, i.e. we have a diagram:

$$\begin{array}{ccc} T & \xrightarrow{\cong} & T' \\ {\scriptstyle\alpha}\downarrow & \swarrow_{g\circ\alpha} & \\ X. & & \end{array} \tag{3.2}$$

If T is a G-torsor over S, and $\alpha : T \to X$ is the G-map, then $g \circ \alpha$ is also a G-torsor over S and there is an isomorphism between T and T'. So $g \in \widetilde{\mathrm{Aut}(X)}$ and the induced map on π_1 is also an isomorphism. So the natural map f is an equivalence.

□

The proof of Theorem 1.1. Let $\mathcal{X}(\mathbf{\Sigma})$ be a toric Deligne-Mumford stack associated to the stacky fan $\mathbf{\Sigma}$. It is a quotient stack $[Z/G]$, where G is an abelian group scheme and affine. So from Lemma 3.2, the natural map

$$f : \mathfrak{G} \to \mathrm{Aut}(\mathcal{X}(\mathbf{\Sigma}))$$

in the introduction is an equivalence as 2-groups. □

4. The weighted projective linear 2-group

In this section we consider the weighted projective stacks. Their automorphism groups are called weighted projective linear 2-groups.

4.1. **The weighted projective stacks.**

Definition 4.1. *Let $Q = (q_0, \cdots, q_n)$ be a $(n+1)$-tuple of positive integers. The weighted projective stack of type Q, $\mathbf{P}(Q) = \mathbf{P}^n_{q_0,\cdots,q_n}$ is defined to be the quotient stack*

$$\mathbf{P}(Q) := [\mathbb{C}^{n+1} \setminus \{0\}/\mathbb{C}^*],$$

where the action is given by $\lambda(x_0, \cdots, x_n) = (\lambda^{q_0} x_0, \cdots, \lambda^{q_n} x_n)$.

Remark 4.2. *(1) The above $\mathbb{C}^*$-action is free iff $q_i = 1$ for every $i = 0, \cdots, n$. In this case this is the projective space $\mathbf{P}^n$. (2) If $gcd(q_0, \cdots, q_n) = d \neq 1$, then $\mathbf{P}^n_{q_0,\cdots,q_n}$ is reduced. Otherwise there exists a gerbe structure over the underlying orbifold.*

Next we construct weighted projective stacks as toric Deligne-Mumford stacks. Let $d = gcd(q_0, \cdots, q_n)$, $a_0 = \frac{q_0}{d}, \cdots, a_n = \frac{q_n}{d}$ and

$$Q_{red} = (a_0, \cdots, a_n).$$

Then $\mathbf{P}(Q_{red})$ is a simplicial toric orbifold. And we can construct a fan Σ in a rank n lattice $\mathbb{Z}^n$. Actually from Fulton [F], we construct the simplicial fan Σ of $\mathbf{P}(Q_{red})$ as follows. Let the fan Σ be generated by vectors $\{v_0, \cdots, v_n\}$ so that $a_0 v_0 + a_1 v_1 + \cdots + a_n v_n = 0$, then the toric variety X_Σ is the coarse moduli space of the weighted projective stack $\mathbf{P}(Q_{red})$.

Remark 4.3. *The paper of Conrads* [Con] *gives a method to compute the lattice vectors $\{v_0, \cdots, v_n\}$ such that they generate a simplicial fan Σ for the corresponding weighted projective stack $\mathbf{P}(Q_{red})$. Then $\mathbf{\Sigma_{red}} = (\mathbb{Z}^n, \Sigma, \overline{\beta})$, where $\overline{\beta} : \mathbb{Z}^{n+1} \longrightarrow \mathbb{Z}^n$ is given by $\{v_0, \cdots, v_n\}$, is a stacky fan and*

$$\mathcal{X}(\mathbf{\Sigma_{red}}) = \mathbf{P}(Q_{red}).$$

Let $N = \mathbb{Z}^n \oplus \mathbb{Z}_d$ and consider the map

$$\beta : \mathbb{Z}^{n+1} \to N \tag{4.1}$$

given by vectors $\{b_0 = (v_0, 1), b_1 = (v_1, 0), \cdots, b_n = (v_n, 0)\} \subset N$. Then

$$\mathbf{\Sigma} = (N, \Sigma, \beta)$$

is a stacky fan in the sense of Borisov- Chen-Smith [BCS]. From the definition of Gale dual in [BCS] we get the following two diagrams:

$$\begin{array}{ccccccccc} 0 & \longrightarrow & \mathbb{Z} & \longrightarrow & \mathbb{Z}^{n+1} & \xrightarrow{\beta} & N & \longrightarrow & 0 \\ & & \downarrow & & \downarrow & & \downarrow & & \\ 0 & \longrightarrow & \mathbb{Z} & \longrightarrow & \mathbb{Z}^{n+1} & \xrightarrow{\overline{\beta}} & \mathbb{Z}^n & \longrightarrow & 0, \end{array} \tag{4.2}$$

and

$$\begin{array}{ccccccccc} 0 & \longrightarrow & \mathbb{Z}^n & \longrightarrow & \mathbb{Z}^{n+1} & \xrightarrow{\overline{\beta}^\vee} & \mathbb{Z} & \longrightarrow & 0 & \longrightarrow & 0 \\ & & \downarrow & & \downarrow & & \downarrow{\scriptstyle [d]} & & \downarrow & & \\ 0 & \longrightarrow & \mathbb{Z}^n & \longrightarrow & \mathbb{Z}^{n+1} & \xrightarrow{\beta^\vee} & \mathbb{Z} & \longrightarrow & \mathbb{Z}_d & \longrightarrow & 0, \end{array} \tag{4.3}$$

where $\overline{\beta}^\vee$ is given by the matrix

$$[a_0, \cdots, a_n]$$

and $\beta^\vee$ is given by the matrix

$$[q_0, \cdots, q_n].$$

Applying $\mathrm{Hom}_{\mathbb{Z}}(-, \mathbb{C}^*)$ to (4.3) yields

$$1 \longrightarrow \mu_d \longrightarrow \mathbb{C}^* \xrightarrow{\alpha} (\mathbb{C}^*)^{n+1} \longrightarrow (\mathbb{C}^*)^n \longrightarrow 1, \tag{4.4}$$

where α is given by the matrix

$$\begin{bmatrix} \lambda^{q_0} \\ \vdots \\ \lambda^{q_n} \end{bmatrix}.$$

From the construction of toric Deligne-Mumford stack in the introduction the weighted projective stack $\mathbf{P}(Q)$ is the toric Deligne-Mumford stack

$$\mathcal{X}(\mathbf{\Sigma}) = [\mathbb{C}^{n+1} \setminus \{0\}/\mathbb{C}^*],$$

where the action is given by $\lambda(x_0, \cdots, x_n) = (\lambda^{q_0} x_0, \cdots, \lambda^{q_n} x_n)$. Thus we have proved the following proposition.

Proposition 4.4. *Every weighted projective stack of type Q is a toric Deligne-Mumford stack $\mathcal{X}(\mathbf{\Sigma})$ with an underlying stacky fan $\mathbf{\Sigma}$.* □

Example 4.5. *For $Q = (4, 6, 8)$, we have $d = 2$ and $Q_{red} = (2, 3, 4)$. The weighted projective stack $\mathbf{P}(Q_{red}) = \mathbf{P}_{2,3,4}$ has a simplicial fan Σ generated by $v_0 = (-3, -2), v_1 = (2, 0), v_2 = (0, 1)$, see* [Jiang]. *We have the stacky fan*

$$\mathbf{\Sigma} = (N, \Sigma, \beta),$$

where $N = \mathbb{Z}^2 \oplus \mathbb{Z}_2$, and $\beta : \mathbb{Z}^3 \to N$ is given by the vectors

$$\{(2, 0, 1), (0, 1, 0)), (-3, -2, 0)\}.$$

The Gale dual map

$$\beta^\vee : \mathbb{Z}^3 \to \mathbb{Z}$$

is given by the matrix $[4, 6, 8]$. So taking Hom$(-, \mathbb{C}^*)$ functor we have that the map

$$\alpha : \mathbb{C}^* \to (\mathbb{C}^*)^3$$

is given by the matrix

$$\begin{bmatrix} \lambda^4 \\ \lambda^6 \\ \lambda^8 \end{bmatrix}.$$

The toric Deligne-Mumford stack $\mathcal{X}(\mathbf{\Sigma}) = [\mathbb{C}^3 \setminus \{0\}/\mathbb{C}^*]$ *is the weighted projective stack* $\mathbf{P}(Q)$ *which is the nontrivial* μ_2*-gerbe over* $\mathbf{P}(Q_{red})$.

4.2. **The weighted projective linear 2-groups.** From Section 4.1 let $Z = \mathbb{C}^{n+1} \setminus \{0\}$, then $\mathrm{Aut}(Z) = GL(n, \mathbb{C})$, the nonsingular $n \times n$ complex matrices over $\mathbb{C}$. Since the group $\mathbb{C}^*$ embeds into the automorphism group $\mathrm{Aut}(Z) = GL(n, \mathbb{C})$ as a diagonal subgroup according to the map α in the exact sequence (4.4), we have the following Lemma.

Lemma 4.6. *The centralizer* $\widetilde{Aut^0(Z)}$ *of* $\mathbb{C}^*$ *in* $Aut(Z)$ *coincides with the normalizer* $\widetilde{Aut(Z)}$ *of* $\mathbb{C}^*$ *in* $Aut(Z)$. $\square$

So we have a natural map $\varphi : \mathbb{C}^* \to \widetilde{\mathrm{Aut}(Z)}$ and from the Introduction we have a 2-group:

$$\mathbf{PGL} = [\mathbb{C}^* \xrightarrow{\varphi} \widetilde{\mathrm{Aut}(Z)}]$$

which is called the weighted projective linear 2-group. From the main result of the paper or the Proposition in [BN], let $\mathrm{Aut}(\mathbf{P}(Q))$ be the automorphism group of the weighted projective stack $\mathbf{P}(Q)$, we have:

Corollary 4.7. *The natural map* $f : \mathbf{PGL} \to Aut(\mathbf{P}(Q))$ *is an isomorphism.* $\square$

Example 4.8. *Let* $B = \mathbf{P}^d$ *be the* d*-dimensional projective space. We give stacky fan* $\mathbf{\Sigma} = (N, \Sigma, \beta)$ *as follows. Let* $N = \mathbb{Z}^d \oplus \mathbb{Z}_r$ *and* $\beta : \mathbb{Z}^{d+1} \longrightarrow N$ *be the map determined by the vectors:*

$$\{(1, 0, \ldots, 0, 0), (0, 1, \ldots, 0, 0), \ldots, (0, 0, \ldots, 1, 0), (-1, -1, \ldots, -1, 1)\}.$$

Then $DG(\beta) = \mathbb{Z}$*, and the Gale dual* $\beta^\vee$ *is given by the matrix*

$$[r, r, \ldots, r].$$

So we have the following exact sequences:

$$0 \longrightarrow \mathbb{Z} \longrightarrow \mathbb{Z}^{d+1} \xrightarrow{\beta} \mathbb{Z}^d \oplus \mathbb{Z}_r \longrightarrow 0 \longrightarrow 0,$$

$$0 \longrightarrow \mathbb{Z}^d \longrightarrow \mathbb{Z}^{d+1} \xrightarrow{\beta^\vee} \mathbb{Z} \longrightarrow \mathbb{Z}_r \longrightarrow 0.$$

Then we obtain the exact sequence:

$$1 \longrightarrow \mu_r \longrightarrow \mathbb{C}^* \xrightarrow{\alpha} (\mathbb{C}^*)^{d+1} \longrightarrow (\mathbb{C}^*)^d \longrightarrow 1. \tag{4.5}$$

The toric Deligne-Mumford stack $\mathcal{X}(\mathbf{\Sigma}) := [\mathbb{C}^{d+1} - \{0\}/\mathbb{C}^*]$ *is the canonical* μ_r*-gerbe over the projective space* $\mathbf{P}^d$ *coming from the canonical line bundle, where the* $\mathbb{C}^*$ *action is given by*

$$\lambda \cdot (z_1, \ldots, z_{d+1}) = (\lambda^r \cdot z_1, \ldots, \lambda^r \cdot z_{d+1}).$$

Denote this toric Deligne-Mumford stack by $\mathcal{G}_r = \mathbf{P}(r, \ldots, r)$*. Let* $Z = \mathbb{C}^{d+1} \backslash \{0\}$*, we have that* $Aut(Z) = GL(d+1, \mathbb{C})$*. The map* α *in (4.5) is given by* $\lambda \longmapsto diag(\lambda^r, \cdots, \lambda^r)$*, so* $\widetilde{Aut(Z)} \cong GL(d+1, \mathbb{C})$*. The automorphism 2-group of* $\mathcal{G}_r$ *is isomorphic to*

$$\mathfrak{G} = [\mathbb{C}^* \xrightarrow{\varphi} GL(d+1, \mathbb{C})]$$

where φ *is given by*

$$\lambda \longmapsto \begin{bmatrix} \lambda^r & & \\ & \vdots & \\ & & \lambda^r \end{bmatrix}.$$

Example 4.9. *We give stacky fan* $\mathbf{\Sigma} = (N, \Sigma, \beta)$ *as follows. Let* $N = \mathbb{Z} \oplus \mathbb{Z}_2$ *and* $\beta : \mathbb{Z}^2 \longrightarrow N$ *be the map determined by the vectors:*

$$\{(2, 0), (-3, 1)\}.$$

Then $DG(\beta) = \mathbb{Z}$*, and the Gale dual* $\beta^\vee$ *is given by the matrix* $[6, 4]$*. So we have the following exact sequences:*

$$0 \longrightarrow \mathbb{Z} \longrightarrow \mathbb{Z}^2 \xrightarrow{\beta} \mathbb{Z} \oplus \mathbb{Z}_2 \longrightarrow 0 \longrightarrow 0,$$

$$0 \longrightarrow \mathbb{Z} \longrightarrow \mathbb{Z}^2 \xrightarrow{\beta^\vee} \mathbb{Z} \longrightarrow \mathbb{Z}_2 \longrightarrow 0.$$

Then we obtain the exact sequence:

$$1 \longrightarrow \mu_2 \longrightarrow \mathbb{C}^* \xrightarrow{\alpha} (\mathbb{C}^*)^2 \longrightarrow (\mathbb{C}^*) \longrightarrow 1. \tag{4.6}$$

The toric Deligne-Mumford stack $\mathcal{X}(\mathbf{\Sigma}) := [\mathbb{C}^2 - \{0\}/\mathbb{C}^*]$ *is the canonical* μ_2*-gerbe over the weighted projective stack* $\mathbf{P}_{2,3}$ *coming from the canonical line bundle, where the* $\mathbb{C}^*$ *action is given by*

$$\lambda \cdot (z_1, z_2) = (\lambda^6 \cdot z_1, \lambda^4 \cdot z_2).$$

The weighted projective stack $\mathbf{P}_{4,6}$ *is the moduli stack* $\overline{\mathcal{M}}_{1,1}$ *of elliptic curves with one marked point. Let* $Z = \mathbb{C}^2 \setminus \{0\}$*, we have that* $Aut(Z) = GL(2, \mathbb{C})$*. The map* α *in (4.6) is given by* $\lambda \longmapsto diag(\lambda^6, \lambda^4)$*, so* $\widetilde{Aut(Z)} \cong \mathbb{C}^* \times \mathbb{C}^*$*. The automorphism 2-group of* $\mathbf{P}_{4,6}$ *is isomorphic to*

$$\mathfrak{G} = [\mathbb{C}^* \xrightarrow{\varphi} \mathbb{C}^* \times \mathbb{C}^*]$$

where φ *is given by*

$$\lambda \longmapsto \begin{bmatrix} \lambda^6 \\ \lambda^4 \end{bmatrix}.$$

References

[BN] K. Behrend, B. Noohi, Uniformization of Deligne-Mumford curves, *J. Reine Angew Math.* 599(2006),111-153, math.AG/0504309.

[BCS] L. Borisov, L. Chen and G. Smith, The orbifold Chow ring of toric Deligne-Mumford stacks, *J. Amer. Math. Soc.* 18 (2005), no.1, 193-215, math.AG/0309229.

[Cox] D. Cox, The homogeneous coordinate ring of a toric variety, *J. of Algebraic Geometry*, 4 (1995), 17-50.

[Con] H. Conrads, Weighted projective spaces and reflexive simplices, *Manuscripta. Mathematica*,Vol 107, No.2, 2002, 215-227.

[F] W. Fulton, *Introduction to toric varieties*, Annals of Mathematics Studies 131, Princeton University Press, Princeton, NJ, 1993.

[Jiang] Y. Jiang, The Chen-Ruan cohomology of weighted projective spaces, to appear in *Canad. J. Math.*, math.AG/0304140.

[KM] S. Keel and S. Mori, Quotients by groupoids, *Ann. of Math.* (2) 145 (1997), no.1, 193-213.

[L-MB] G. Laumon and L. Moret-Bailly, Champs algébriques. (French) [Algebraic stacks] Ergebnisse der Mathematik und ihrer Grenzgebiete. 3. Folge. A Series of Modern Surveys in Mathematics [Results in Mathematics and Related Areas. 3rd Series. A Series of Modern Surveys in Mathematics], 39. Springer-Verlag, Berlin, 2000.

[Noohi] B. Noohi, Notes on 2-groupoids, 2-groups and crossed-modules, math.CT/0512106.

[V] A. Vistoli, Intersection theory on algebraic stacks and on their moduli spaces, *Invent. Math.*, 97 (1989) 613-670.

Department of Mathematics, University of British Columbia, 1984 Mathematics Road, Vancouver, BC V6T 1Z2, Canada

E-mail address: jiangyf@math.ubc.ca

Current Address: Department of Mathematics, University of Utah, 155 S 1400 E JWB 233, Salt Lake city, UT 84112, USA